



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> :  
C07K 15/00, C08G 69/10  
C07K 13/00, 7/00, A61K 37/02  
A61K 37/24, C07D 263/44

A1

(11) International Publication Number:

WO 91/06569

(43) International Publication Date:

16 May 1991 (16.05.91)

(21) International Application Number: PCT/US90/06213

(22) International Filing Date: 31 October 1990 (31.10.90)

(30) Priority data:

430,990

3 November 1989 (03.11.89) US

(60) Parent Application or Grant

(63) Related by Continuation

US

430,990 (CIP)

Filed on

3 November 1989 (03.11.89)

(71) Applicant (for all designated States except US): BOOTS  
PHARMACEUTICALS INC. [US/US]; 300 Tri State  
International Centre, Lincolnshire, IL 60069-4415 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only) : LATHAM, Keith, Roger [US/US]; 5600 Foggy Lane, Rockville, MD 20855 (US). LATHAM, Vincent, Ernest [US/US]; 608 Rainbow Boulevard, Lady Lake, FL 32159 (US).

(74) Agent: MURRAY, Robert, B.; Armstrong, Nikaido, Marmelstein, Kubovcik & Murray, Suite 1000, 1725 K Street, N.W., Washington, DC 20006 (US).



(81) Designated States: AT, AT (European patent), AU, BE (European patent), BG, BR, CA, CH, CH (European patent), DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GB, GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU, LU (European patent), NL, NL (European patent), NO, RO, SE, SE (European patent), SU, US.

## Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: IODOTHYRONINE POLYMERS



## (57) Abstract

Iodothyronine polymers having a plurality of recurring units of formula (I), in which A is iodo and B, C, and D are independently H or iodo are described. Polymers in which A and C are iodo and B and D are independently H or iodo and in which substantially all of the recurring units are L-stereoisomers, have utility in treating thyroid hormone deficiencies. The polymers are preferably prepared by polymerization of N-carboxyanhydrides of formula (III).

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CI	Côte d'Ivoire	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

- 1 -


IODOTHYRONINE POLYMERS

This invention relates to iodothyronine polymers which have utility in the treatment of thyroid hormone deficiencies, to pharmaceutical compositions containing  
5 iodothyronine polymers and to the use of iodothyronine polymers in the treatment of thyroid hormone deficiencies. The iodothyronine polymers of the present invention contain recurring units linked by -NHCO- bridging groups and are therefore polypeptides.

10 Thyroid hormone deficiencies are disease states in which insufficient thyroid hormone is released in the body causing a slowing down of all the metabolic processes of the body and, in children, causing poor mental and physical development. Dessicated thyroid  
15 glands obtained from the ox, sheep or pig have been used for many years to treat thyroid hormone deficiencies. However the actual thyroid hormone dose from dessicated thyroid glands is difficult to regulate due to variations in iodine content between  
20 preparations. More recently synthetic levothyroxine (LT<sub>4</sub>) has been used to treat thyroid hormone deficiencies.

The present invention provides a substantially pure synthetic iodothyronine polymer having a plurality  
25 of recurring units, which may be the same or different, of formula I

- 2 -



in which A is iodo and B, C and D are independently H or iodo. In preferred iodothyronine polymers substantially all of the recurring units of formula I are in the same stereoisomeric form. In particularly preferred iodothyronine polymers substantially all of the recurring units of formula I are L-stereoisomers. In preferred iodothyronine polymers A and C are iodo and B and D are independently H or iodo. In particularly preferred iodothyronine polymers A and C are iodo and at least one of B and D is iodo. The average number of the recurring units may vary from about 5 to about 400, preferably from about 10 to about 400, more preferably from about 20 to about 200, or from about 30 to about 150 or from about 80 to about 120.

The recurring units of formula I are derivatives of one or more iodothyronine compounds selected from the group consisting of 3-T<sub>1</sub>, 3,3'-T<sub>2</sub>, 3,5-T<sub>2</sub>, rT<sub>3</sub>, T<sub>3</sub> and T<sub>4</sub> as defined in Table I below.

- 3 -

Table I

	3,3',5,5'-Tetraiodothyronine	T <sub>4</sub>
	3,3',5-Triiodothyronine	T <sub>3</sub>
	3,3',5'-Triiodothyronine	rT <sub>3</sub>
5	3,5-Diiodothyronine	3,5-T <sub>2</sub>
	3,3'-Diiodothyronine	3,3'-T <sub>2</sub>
	3-Monoiodothyronine	3-T <sub>1</sub>

When the iodothyronine polymers of the present invention are used to treat thyroid hormone deficiencies, substantially all of the recurring units of formula I are the physiologic L-stereoisomer. That is, at least 90%, preferably 95%, and most preferably greater than 99% of the recurring units are the physiologic L-stereoisomer. In the iodothyronine polymers used to treat thyroid hormone deficiencies, the recurring units of formula I are derivatives of the pharmacologically active iodothyronine compounds identified in Table II.

Table II

20	3,3',5,5'-Tetraiodo-L-thyronine	Thyroxine	LT <sub>4</sub>
	3,3',5-Triiodo-L-thyronine	Liothyronine	LT <sub>3</sub>
	3,3',5'-Triiodo-L-thyronine	Reverse T <sub>3</sub>	rLT <sub>3</sub>
	3,3'-Diiodo-L-thyronine	-	3,3'-LT <sub>2</sub>

25 In one embodiment of the invention the iodothyronine polymer is a homopolymer as defined below in Table III.

- 4 -

Table III

	<u>Abbreviation</u>	<u>Substitution in Formula I</u>			
		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>
	poly-T <sub>4</sub>	I	I	I	I
5	poly-T <sub>3</sub>	I	I	I	H
	poly-rT <sub>3</sub>	I	H	I	I
	poly-3,5-T <sub>2</sub>	I	I	H	H
	poly-3,3'-T <sub>2</sub>	I	H	I	H
	poly-3-T <sub>1</sub>	I	H	H	H

- 10 In each such homopolymer, substantially all of the recurring units will be those identified above. That is, at least 90%, preferably at least 95%, and most preferably at least 99% of the recurring units of each homopolymer will contain the substituents identified in
- 15 Table III. The iodothyronine homopolymers in which substantially all of the recurring units of formula I are L-stereoisomers and in which A, B, C and D are iodo, or in which A, B and C are iodo and D is H have utility in the treatment of thyroid hormone deficiency.
- 20 The preferred homopolymers for use in therapy are poly-LT<sub>4</sub> and poly-LT<sub>3</sub>.

The iodothyronine polymers of the present invention in which substantially all of the recurring units of formula I are L-stereoisomers and A and C are

25 iodo and B and D are independently H or iodo have utility in the treatment of thyroid hormone deficiencies in human and other mammals. LT<sub>4</sub> is the primary thyroid hormone in mammals, but LT<sub>3</sub> is also released by the thyroid gland and is also active as a thyroid hormone. LT<sub>4</sub> and LT<sub>3</sub> are found in the blood in


30 an approximate 4:1 ratio. In one embodiment of the present invention, the iodo-thyronine polymer contains recurring units derived from both LT<sub>4</sub> and LT<sub>3</sub> to form a

- 5 -

copolymer of these units. Preferably, from about 70 to about 90% of the recurring units are the  $LT_4$  derivatives and about 10 to about 30% of the recurring units are the  $LT_3$  derivatives. Most preferably, the ratio is approximately 4:1 i.e. about 80% of the recurring units are  $LT_4$  and approximately 20% of such units are  $LT_3$ . This copolymer is referred to herein as poly- $LT_4/LT_3$ .

It has been found that  $rLT_3$  may also have a role in thyroid hormone function in humans and other mammals. A further aspect of the present invention provides an iodothyronine polymer in which a fraction of the recurring units are derived from  $rLT_3$ . In such an iodothyronine polymer, approximately 70-89% of the recurring units are derived from  $LT_4$ , approximately 10-29% of the recurring units are derived from  $LT_3$ , and the remainder of the recurring units are derived from  $rLT_3$ . Most preferably, approximately 80% of such units are derived from  $LT_4$ , approximately 15% of such units are derived from  $LT_3$ , and approximately 5% of such units are derived from  $rLT_3$ . This heteropolymer is referred to herein as poly- $LT_4/LT_3/rLT_3$ .

A further aspect of the present invention provides iodothyronine polymers which contain further recurring units of formula II



in which R is a residue of any of the amino acids commonly found in nature. These amino acids are listed by Lehninger in Principles of Biochemistry (1982) published by Worth Publishers Inc of New York (see page 96) as alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine,

- 6 -

isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine. The properties of the iodothyronine polymers are modified by copolymerization with these amino acids. For example, if greater aqueous solubility is desired, hydrophilic amino acids, like lysine, arginine, aspartic acid, glutamic acid, serine or threonine can be copolymerized into the polypeptide chain. The amount of amino acid incorporated will be dependent upon the particular iodothyronine/amino acid copolymer(s), and the properties desired. The recurring units of formula II may comprise up to 50%, preferably up to 66%, most preferably up to 80% of the recurring units in the copolymers.

Iodothyronine polymers of the present invention in which substantially all of the recurring units of formula I are the L-stereoisomers and in which the recurring units are of formula I in which A and C are iodo and B and D are independently H or iodo have utility in the treatment of thyroid hormone deficiencies. A further aspect of the present invention therefore provides a pharmaceutical composition suitable for treating thyroid hormone deficiencies which comprises a pharmaceutically acceptable diluent or carrier and a pharmacologically active ingredient consisting of a pharmaceutically effective amount of a substantially pure synthetic iodothyronine polymer having a plurality of recurring units, which may be the same or different, of the formula I in which A and C are iodo and B and D are independently H or iodo, in which polymer substantially all of the recurring units of formula I are L-stereoisomers. Preferably at least one of B or D is iodo. The average number of the recurring units may vary from about 5 to about 400, preferably from about 10 to about 400, more preferably from about 20 to about



- 7 -

200, or from about 30 to about 150 or from about 80 to about 120. The iodothyronine polymer in these pharmaceutical compositions may be poly-LT<sub>4</sub>, poly-LT<sub>3</sub>, poly-LT<sub>4</sub>/LT<sub>3</sub>, poly-LT<sub>4</sub>/LT<sub>3</sub>/rLT<sub>3</sub> or mixtures thereof. Particularly preferred compositions are those in which the iodothyronine polymer is poly-LT<sub>4</sub>, poly-LT<sub>3</sub>, a mixture of poly-LT<sub>4</sub> and poly-LT<sub>3</sub> in which the ratio of poly-LT<sub>4</sub> to poly-LT<sub>3</sub> lies in the range 7:3 to 9:1 preferably about 4:1 or a mixture of poly-LT<sub>4</sub>, poly-LT<sub>3</sub> and poly-rLT<sub>3</sub> in which the ratio of poly-LT<sub>4</sub> to poly-LT<sub>3</sub> and poly-rLT<sub>3</sub> is approximately 80:15:5.

In therapeutic use, the iodothyronine polymer is preferably administered orally. Thus the therapeutic compositions of the present invention may take the form of any of the known pharmaceutical compositions for oral administration. Pharmaceutically acceptable carriers are well known in the art of pharmacy. The compositions of the invention may contain 0.1-90% by weight of iodothyronine polymer. The compositions of the invention are generally prepared in unit dosage form.

Compositions for oral administration are the known pharmaceutical forms for such administration, for example tablets, capsules, syrups and aqueous or oily suspensions. The diluent or carrier used in the preparation of these compositions can be any of the materials known in the pharmacists' art. Tablets may be prepared by mixing the iodothyronine polymer with an inert diluent such as calcium phosphate in the presence of disintegrating agents, for example maize starch, and lubricating agents, for example magnesium stearate, and tableting the mixture by known methods. The tablets may be formulated in a manner known to those skilled in the art so as to give a sustained release of the compounds of the present invention. Such tablets may,

- 8 -

if desired, be provided with enteric coatings by known methods, for example by the use of cellulose acetate phthalate. Similarly, capsules, for example hard or soft gelatin capsules, containing the active compound with or without added excipients, may be prepared by conventional means and, if desired, provided with enteric coatings in a known manner. The tablets and capsules may conveniently each contain 12.5 to 500 microgrammes of the iodothyronine polymer. Other compositions for oral administration include, for example, aqueous suspensions containing the iodothyronine polymer in an aqueous medium in the presence of a non-toxic suspending agent such as sodium carboxymethylcellulose, and oily suspensions containing a compound of the present invention in a suitable vegetable oil, for example arachis oil.

An alternative route of administration of the iodothyronine polymer is by means of an implant. In this case a pellet containing iodothyronine polymers containing large numbers of recurring units is implanted under the skin of the patient and pharmacologically active amounts of the iodothyronine polymer are then released over an extended period of time, suitably over several weeks or months.

In some formulations it may be beneficial to use the compounds of the present invention in the form of particles of very small size, for example as obtained by fluid energy milling.


The pharmaceutical compositions containing a therapeutically effective amount of the iodothyronine polymers of the present invention in which substantially all of the recurring units of formula I are the L-stereoisomers may be used to treat thyroid hormone deficiencies in warm blooded animals including

- 9 -

human beings. In such treatment the amount of the iodothyronine polymer administered per day is under the control of the prescribing physician and will depend inter alia on the age of the patient and on the severity of the condition to be treated but will usually lie in the range 12.5 to 1000 micrograms per day, preferably 25 to 400 micro-grams per day, most preferably 50 to 300 micrograms per day given in single or divided doses at one or more times during the day.

In the case of iodothyronine polymers containing recurring units of formula II, the amount of the polymer administered may be higher than quoted above but will be such as to give rise to equivalent amounts of the pharmacologically active iodothyronine moieties.

A preferred method of synthesizing the iodothyronine polymers of the present invention comprises the polymerisation of one or more N-carboxyanhydrides (NCA) of formula III



in which A is iodo, and B, C, and D are H or iodo. Suitable N-carboxyanhydrides of formula III are identified in Table IV.


Table IV

	<u>Abbreviation</u>	<u>Substitution in Formula III</u>			
		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>
	T <sub>4</sub> -NCA	I	I	I	I
5	T <sub>3</sub> -NCA	I	I	I	H
	rT <sub>3</sub> -NCA	I	H	I	I
	3,5-T <sub>2</sub> -NCA	I	I	H	H
	3,3'-T <sub>2</sub> -NCA	I	H	I	H
	3-T <sub>1</sub> -NCA	I	H	H	H

10        When the desired iodothyronine polymer is one in which substantially all of the recurring units are in the same stereoisomeric form (preferably the L-stereoisomeric form), the N-carboxyanhydride of formula III should be substantially all in that same  
15 stereoisomeric form.

When the desired iodothyronine polymer is a homopolymer, one N-carboxyanhydride of formula III is used. However, if the desired iodothyronine polymer is a copolymer or heteropolymer, then two or more  
20 N-carboxyanhydrides of formula III are used in the same molar proportions as is desired in the iodothyronine polymer. For example, when the desired iodothyronine polymer is a copolymer of LT<sub>4</sub> and LT<sub>3</sub>, the N-carboxyanhydrides of LT<sub>4</sub> and of LT<sub>3</sub> are used in the molar  
25 ratio desired in the iodothyronine copolymer and when the desired iodothyronine polymer is a heteropolymer of LT<sub>4</sub>, LT<sub>3</sub> and rLT<sub>3</sub>, the N-carboxyanhydrides of LT<sub>4</sub>, LT<sub>3</sub> and rLT<sub>3</sub> are used in the molar ratio desired in the final copolymer. When the iodothyronine polymer also  
30 contains one or more further recurring units of formula II, an N-carboxyanhydride of formula IV

- 11 -



may be used in addition to the N-carboxyanhydride of formula III. The N-carboxyanhydrides of formula III and formula IV are used in the molar ratio desired in the final iodothyronine polymer

5       The polymerization preferably comprises reacting one or more compounds of formula III in an anhydrous solvent, preferably at a concentration of about 5-40% and a temperature from about 0°C to the boiling point of the solvent for sufficient time to complete the  
10       polymerisation. The reaction is continued until polymerisation is complete as indicated, for example, by product precipitation, cessation of CO<sub>2</sub> evolution, the attainment of maximum viscosity or the absence of the starting material, as indicated for example, by  
15       spectroscopic examination of the reaction mixture. Examples of suitable anhydrous solvents include ethers, such as dioxane or tetrahydrofuran, aromatic solvents, such as benzene, chlorobenzene, and toluene, and other solvents, such as dimethyl formamide, ethyl acetate and  
20       dimethyl sulphoxide. A preferred solvent is dioxane or tetrahydrofuran.

Preferably, a base is used as an initiator in the above reaction. An inorganic or organic base may be used, although an organic base is preferred. Examples  
25       of suitable bases include organic amines such as n-butylamine, triethylamine, tributylamine, triamylamine, diisopropylethylamine or alkali metal alkoxides such as sodium methoxide or sodium ethoxide. The molar

- 12 -

ratio of the N-carboxyanhydride derivative of formula III to the initiator lies in the range 20 to 400 preferably 30 to 150 more preferably 50 to 100. Most preferably, the base is sodium methoxide.

5           The iodothyronine polymers of the present invention may be prepared by the further iodination of iodothyronine polymers having recurring units of formula I in which A is iodo, B is H or iodo and C and D are H. These iodothyronine polymer starting  
10 materials are identified in Table III as poly-3,5-T<sub>2</sub> and poly-3-T<sub>1</sub> respectively. If these polymers are subjected to vigorous iodination conditions, for example by the use of excess potassium triiodide or of iodine monochloride as the iodinating agent,  
15 iodothyronine polymers can be produced in which both C and D are iodo. Thus poly-3,5-LT<sub>2</sub> gives poly-LT<sub>4</sub> and poly-3-LT<sub>1</sub> gives poly-rLT<sub>3</sub>. By using less vigorous iodination conditions, for example potassium triiodide in diethylamine, it is possible to produce  
20 iodothyronine polymer in which C is iodo and D is H. Thus poly-3,5-LT<sub>2</sub> would give poly-LT<sub>3</sub> and poly-3-LT<sub>1</sub> would give poly-3,3'-LT<sub>2</sub>. By choosing iodination conditions in which iodination is incomplete, for example by restricting the amount of iodinating agent,  
25 it is possible to produce iodothyronine polymers in which C is iodo and in which D is iodo in only some of the recurring units. For example, copolymers of LT<sub>4</sub> and LT<sub>3</sub> may be prepared by partial iodination of poly-3,5-LT<sub>2</sub>.


30           The iodothyronine polymers of the present invention may also be prepared by the condensation of the amino acids from which the recurring units of formula I are derived in the presence of a dehydrating agent such as dicyclohexylcarbodiimide which is  
35 converted into dicyclohexylurea when it removes the

- 13 -

elements of water from two amino acid residues to form a peptide bond between them. Copolymers and heteropolymers may be prepared by condensing mixtures of two or more amino acids.

5 Homogeneous polymerizations to form heteropolymers result in a random distribution of each component in the polymer. However, when sequence-specific copolymer or heteropolymer combinations are  
10 is useful. The preferred method of synthesis is through the t-BOC or f-MOC intermediates of the iodothyronines and other amino acids if applicable. Synthesis proceeds as previously described in Groginski, Amer. Biotech. Lab., May/June:38-51 (1986),  
15 incorporated herein by reference.

The N-carboxyanhydride of formula III may be prepared by reacting the appropriate iodothyronine with a carbonylating reagent to form the N-carboxyanhydride. More particularly, they are synthesized by reacting a  
20 compound represented by the formula V



or a salt thereof with a carbonylating reagent and recovering the reaction product. Preferably, the

- 14 -

reaction takes place in the presence of an anhydrous solvent, such as tetrahydrofuran (THF), but other appropriate solvents known to those skilled in the art can be used. A preferred carbonylating reagent is  
5 hexachlorodimethylcarbonate,  $(\text{CCl}_3\text{O})_2\text{CO}$ . This normally solid chemical is commercially available from Aldrich Chemicals under the trade name Triphosgene. Typically, the iodothyronine is suspended in the anhydrous solvent and the hexachlorodimethylcarbonate is added. An  
10 excess of the hexachlorodimethylcarbonate is used. Alternatively, phosgene gas may be used as the carbonylating agent.

After it has been prepared the iodothyronine polymer may be precipitated and the resulting solid  
15 collected by filtration and may be further purified, if necessary, by recrystallisation. The product is advantageously dried by lyophilisation. In this process the precipitated iodothyronine polymer, if  
20 necessary following recrystallisation, is prefrozen or frozen in situ by evaporative cooling in vacuo with sufficient external heat provided to obtain a product with the desired, preferably less than 0.1%, moisture content.

The N-carboxyanhydrides of formula III obtained by  
25 the above synthesis are typically recrystallised to obtain intermediates of pharmaceutical purity. Since the prepared anhydrides are moisture sensitive, special care to ensure anhydrous conditions of the reaction, recovery and storage of the N-carboxyanhydrides is  
30 important. This synthetic procedure may also be used to prepare the N-carboxyanhydride derivatives of formula IV for copolymerization or heteropolymerisation with the N-carboxyanhydride derivatives of formula III to prepare iodothyronine polymers having further  
35 recurring units of formula II.



- 15 -

Compounds of formula V are prepared by methods which are well known in the art such as those described, for example, in US Patents 2579668, 2886592, 2889363, 2889364, 3477954 and 3577535.

5           The iodothyronine polymers of the present invention provide a method of delivering thyroid hormones to a patient in need thereof. Because the thyroid hormones are released by digestive proteolysis of the iodothyronine polymers of this invention, it is  
10       expected that the use of the iodothyronine polymers would have a long physiologic effect because of the sustained release from the polymers of the monomeric thyroid hormones. The use of copolymers containing recurring units derived from two or more thyroid  
15       hormones or mixtures of homopolymers of the thyroid hormones in the appropriate ratio provides a means whereby the naturally occurring ratio of thyroid hormones may be duplicated. Attempts have been made before to duplicate this naturally occurring ratio by  
20       administering a mixture of  $LT_3$  and  $LT_4$  (see for example US Patents 3477954 and 3577535). However,  $LT_3$  has a short half life in the blood compared to the half life of  $LT_4$  and so the desired ratio cannot be maintained over a long period of time. The iodothyronine polymers  
25       of the present invention are solid materials which may be readily handled and formulated to give stable, consistent pharmaceutical compositions for the treatment of thyroid hormone deficiencies.

30           The invention will now be illustrated by the following Examples.

Example 1

Hexachlorodimethylcarbonate (10 g) was added to a suspension of 3,5-diiodo-L-thyronine (26.3) in anhydrous tetrahydrofuran (125 ml) and the mixture was heated to 67°C for 15 minutes. Anhydrous tetrahydrofuran (500 ml) and then anhydrous hexane (3000 ml) were added and the mixture stored at 20°C for three hours. 3,5-Diiodo-L-thyronine N-carboxyanhydride (3,5-LT<sub>2</sub>-NCA) was collected by filtration. Yield 24 g.

Example 2

A solution of 3,5-diiodo-L-thyronine N-carboxyanhydride (1 g) in dioxane (10 ml) was stirred rapidly and a 1% solution of sodium methoxide in methanol (0.05 ml) was added. The mixture was stirred for six days at ambient temperature and then petroleum ether (20 ml) was added and the resulting slurry triturated with petroleum ether. Polymeric 3,5-diiodo-L-thyronine (poly 3,5-LT<sub>2</sub>) was collected by filtration and dried in vacuo. Yield 0.96 g.

Example 3

In a similar manner to that described in Example 1, 3,3',5-triiodo-L-thyronine N-carboxyanhydride (LT<sub>3</sub>-NCA) was prepared in 65% yield and polymerised to give polymeric 3,3',5-triiodo-L-thyronine (poly-LT<sub>3</sub>) in 62% yield in a similar manner to that described in Example 2.

Example 4

In a similar manner to that described in Example 1, 3,3',5,5'-tetraiodo-L-thyronine (LT<sub>4</sub>-NCA) was prepared in 22% yield from 3,3',5,5'-tetraiodo-

- 17 -

L-thyronine which had been dried in vacuo at 100°C for 17 hours and then polymerised to give polymeric 3,3',5,5'-tetraiodo-L-thyronine (poly-LT<sub>4</sub>) in 82% yield in a similar manner to that described in Example 2.

#### Example 5

In a similar manner to that described in Example 1, 3,5,5'-triiodo-L-thyronine N-carboxyanhydride (rLT<sub>3</sub>-NCA) was prepared in 51% yield and polymerised to give polymeric 3,5,5'-triiodo-L-thyronine (poly-rLT<sub>3</sub>) in 70% yield in a similar manner to that described in Example 2.

#### Example 6

A mixture of 4 parts of 3,3',5,5'-tetraiodo-L-thyronine N-carboxyanhydride (LT<sub>4</sub>-NCA) prepared in a similar manner to that described in Example 1 and 1 part of 3,3',5-triiodo-L-thyronine N-carboxyanhydride (LT<sub>3</sub>-NCA) prepared in a similar manner to that described in Example 3 was polymerised in a similar manner to that described in Example 2 to give a polymer containing recurring units derived from both LT<sub>3</sub> and LT<sub>4</sub> (poly LT<sub>4</sub>/LT<sub>3</sub>) in 62% yield.

#### Example 7

A mixture of 80 parts 3,3',5,5'-tetraiodo-L-thyronine N-carboxyanhydride (LT<sub>4</sub>-NCA) prepared in a similar manner to that described in Example 4, 15 parts of 3,3',5-triiodo-L-thyronine N-carboxyanhydride (LT<sub>3</sub>-NCA) prepared in a similar manner to that described in Example 3 and 5 parts of 3,5,5'-triiodo-L-thyronine N-carboxyanhydride (rLT<sub>3</sub>-NCA) prepared in a similar manner to that described in Example 5 was

- 18 -

polymerised in a similar manner to that described in Example 2 to give a heteropolymer containing recurring units derived from  $LT_4$ ,  $LT_3$  and  $rLT_3$  (poly  $LT_4/LT_3/-rLT_3$ ). The product was precipitated from the reaction mixture by addition of petroleum ether (2 volumes) and the precipitate dried in vacuo. Yield 55%.

#### Example 8

Polymeric 3,5-diiodo-L-thyronine (poly-3,5- $LT_2$  - 22 g prepared in a similar manner to that described in Example 2) was dissolved in a 33% aqueous solution of diethylamine at 16 to 22°C. A 1.9N iodine solution in concentrated aqueous potassium iodide solution (88 ml) was added with stirring. The mixture was stirred at 4 to 10°C for 2 hours. A precipitate formed which was collected by filtration and washed with water. The filter cake was dissolved in a mixture of ethanol (25 ml) and 2N aqueous sodium hydroxide solution (100 ml). 2N Hydrochloric acid was added to neutralise the solution and a precipitate formed which was collected by filtration and washed with water. The wet filter cake was placed in a freeze drying chamber in vacuo (less than 0.3 mm Hg) to freeze the cake via evaporative cooling. Sufficient heat was provided (shelf temperature 40°C) to reduce the moisture content to less than 0.1% in 24 hours. The product was polymeric 3,3',5,5'-tetraiodo-L-thyronine. Yield 27 g.

#### Example 9

Polymeric 3,5-diiodo-L-thyronine (poly-3,5- $LT_2$  - 18.4 g prepared in a similar manner to that described in Example 2) was dissolved in 33% aqueous diethylamine (185 ml). A solution of potassium triiodide [82 ml of a solution prepared from iodine (26.2 g), potassium iodide (67.8 g) and water (90 ml)] was added with

- 19 -

stirring over 30 minutes. Stirring was continued for 15 minutes and water (111 ml) and then 2N hydrochloric acid were added to cause precipitation. The brown precipitate was collected by filtration and the filter cake dissolved in a mixture of ethanol (1332 ml) and 1N aqueous sodium hydroxide solutions (111 ml). The solution was filtered, heated to boiling and treated with 30% aqueous acetic acid until precipitation commenced. The mixture was cooled in ice and the precipitate collected by filtration and washed with water. The wet filter cake was placed in a freeze drying chamber in vacuo (less than 0.3 mm Hg) to freeze the cake via evaporative cooling. Sufficient heat was provided (shelf temperature 40°C) to reduce the moisture content to less than 0.1% in 24 hours. (Yield 24.6 g).

The product was hydrolysed with acid and high performance liquid chromatography (HPLC) of the hydrolysed product showed the presence of 83.3% LT<sub>4</sub>, 16.4% LT<sub>3</sub> and less than 0.1% of 3,5-LT<sub>2</sub> indicating that the product was a polymer containing recurring units derived from LT<sub>4</sub> and LT<sub>3</sub> (poly LT<sub>4</sub>/LT<sub>3</sub>).

#### Example 10

In a similar manner to that described in Example 8, polymeric 3-iodo-L-thyronine (poly-3LT<sub>1</sub>) was diiodinated to give polymeric 3,5,5'-triiodo-L-thyronine (poly-rLT<sub>3</sub>) in 96% yield. Polymeric 3-iodo-L-thyronine was prepared in a similar manner to that described in Example 2 by polymerisation of 3-iodo-L-thyronine N-carboxyanhydride (3-LT<sub>1</sub>-NCA) which was prepared in 93% yield in a similar manner to that described in Example 1.

- 20 -

Example 11

Direct co-polymerization of  $LT_4$  and glycine was effected by suspending 3,3',5,5'-tetraiodo-L-thyronine (3.88 g) and glycine (Zwitterionic forms) (0.37 g) in anhydrous dioxane (50 ml). Dicyclohexylcarbodiimide (2.47 g) was added with stirring and the reaction was stirred for 4 days at 17-22°C. Acetic acid (0.2 ml) was added to decompose excess dicyclohexylcarbodiimide and petroleum ether (50 ml) was added to precipitate the polymeric product. The product was collected by filtration and the filter cake triturated with ethanol. The polymeric product was dried in vacuo. Yield 5.22 g. HPLC analysis of the acid hydrolysis of the product showed a 50/50 mixture of  $LT_4$  and glycine indicating that the polymeric product was a copolymer containing equimolar amounts of  $LT_4$  and glycine.

Example 12

Boc-protected  $LT_4$  and glycine are prepared as described in Tam et al., Int. J. Peptide Protein Res., 21:57 (1983), and polymerized by standard sequential additions on polystyrene beads as an immobilized support. The following coupling sequence is used as described in Spatola, Amer. Biotech Lab., Dec. 14-22 (1984).

- 21 -

	Stage	Reagent	Repeat	Time
	1	Methylene chloride	5	1
	2	TFA	1	5
5	3	TFA	1	25
	4	Methylene chloride	4	1
	5	Diisopropylethylamine	2	2
	6	Methylene chloride	3	1
	7	DMF	3	1
10	8	Boc-glycine	-	1
	9	DCC	-	-
	10	DMF	1	1
	11	Methylene chloride	3	1
	12	Ethanol	3	1
15	13	DMF	3	1
	14	Boc-LT <sub>4</sub>	-	1
	15	DCC	-	-
	16	DMF	3	1
	17	Methylene chloride	3	1
20	18	Ethanol	3	1

Steps 7 through 18 are repeated 25 or more times to obtain a polymer of 50 or more residues in length. Finally, the copolymer is isolated by cleavage from the resin using HF. The copolymer formed by this reaction has alternating recurring units derived from LT<sub>4</sub> and glycine.

### Example 13

Hexachlorodimethylcarbonate (20 g) was added to a suspension of the sodium salt of 3,5-diiodo-L-thyronine (52.6 g) which had been dried at 100°C in vacuo for 24

- 22 -

hours in anhydrous tetrahydrofuran (250 ml) and the mixture was heated to 65°C. The reaction mixture was allowed to react for five minutes and the solvent was then removed by evaporation. Anhydrous ethyl acetate  
5 (50 ml) and then anhydrous dichloromethane (140 ml) were added and the mixture cooled at 4°C for one hour. 3,5-Diodo-L-thyronine N-carboxyanhydride (3,5-LT<sub>2</sub>-NCA) was collected by filtration, washed with anhydrous hexane and dried in vacuo without heating. Yield  
10 37.6 g.

#### Example 14

A solution of 3,5-diodo-L-thyronine N-carboxyanhydride (37.2 g) in ethyl acetate (372 ml which had been dried over potassium carbonate) was stirred  
15 rapidly and a 1% solution of sodium methoxide in methanol (1.86 ml) was added. The mixture was stirred for four days at ambient temperature and then polymeric 3,5-diodo-L-thyronine (poly 3,5-LT<sub>2</sub>) was collected by filtration, washed with ethyl acetate and  
20 then hexane and dried in vacuo. Yield 24.9 g.

#### Example 15

Polymeric 3,5-diodo-L-thyronine (0.715 g) was ground to a fine powder and dissolved with gentle warming in a mixture of dimethylformamide (8.6 ml),  
25 water (5.7 ml) and diethylamine (4.3 ml) and the solution was cooled to 5-10°C. A solution of potassium triiodide was prepared from iodine (31.5 g) and a 40% solution of potassium iodide in water (100 ml of solution). Four portions of the resulting solution  
30 (0.8 ml each) were added over 30 minutes. The reaction mixture was stirred overnight as the temperature rose to ambient and then poured into acetone (200 ml). A solid was collected by filtration and washed with



- 23 -

degassed water. The washed solid was frozen on dry ice and lyophilised under vacuum (less than 0.3 mm Hg) using a shelf temperature of 35°C to give polymeric 3,3',5-triiodo-L-thyronine (poly-LT<sub>3</sub>). Yield 0.9 g.

5    Example 16

Polymeric 3,5-diiodo-L-thyronine (1 g) was ground to a fine powder and dissolved with gentle warming in dimethylformamide (5 ml). A mixture of glacial acetic acid (2 ml) and iodine monochloride (0.8 g) was added  
10 over twenty minutes with rapid mixing and the resulting mixture heated to 60°C. Glacial acetic acid (5 ml) and then water (12 ml) were added dropwise and the mixture was reheated to 60°C. Potassium bisulphite (0.3 g) was added and the mixture cooled in ice. The resulting  
15 precipitate was collected by filtration and washed with water. The washed solid was frozen on dry ice and lyophilised under vacuum (less than 0.3 mm Hg) using a shelf temperature of 35°C to give polymeric 3,3',5,5'-tetraiodo-L-thyronine (poly-LT<sub>4</sub>). Yield 1.56 g.

20    Example 17

Hexachlorodimethylcarbonate (2 g) was added to a suspension of the sodium salt of 3,3',5-triiodo-L-thyronine (6.73 g) which had been dried at 100°C in vacuo for 24 hours in anhydrous tetrahydrofuran (25 ml)  
25 and the mixture was heated to 65°C. The reaction mixture was allowed to react for ten minutes and the solvent was then removed by evaporation. Anhydrous ethyl acetate (7 ml) was added to dissolve the residue and then anhydrous dichloromethane (20 ml) was added  
30 and the mixture cooled at 4°C for one hour. 3,3',5-Triiodo-L-thyronine N-carboxyanhydride (LT<sub>3</sub>-NCA) was collected by filtration, washed with anhydrous

- 24 -

hexane and dried in vacuo without heating. Yield 2.8 g.

Example 18

A solution of 3,3',5-triiodo-L-thyronine  
5 N-carboxyanhydride (1.0 g) in anhydrous dioxane (10 ml)  
was stirred rapidly and a 1% solution of sodium  
methoxide in methanol (0.05 ml) was added. The mixture  
was stirred for four days at ambient temperature and  
then water (10 ml) was added with vigorous stirring.  
10 Polymeric 3,3',5-triiodo-L- thyronine (poly-LT<sub>3</sub>) was  
collected by filtration and washed with water. The  
washed solid was frozen on dry ice and lyophilised  
under vacuum (less than 0.3 mm Hg) using a shelf  
temperature of 35°C. Yield 0.68 g.

15 Example 19

A suspension of the sodium salt of 3,3',5,5'-  
tetraiodo-L-thyronine (1.55 g) which had been dried at  
100°C in vacuo for 24 hours in anhydrous tetra-  
hydrofuran was cooled and hexachlorodimethylcarbonate  
20 (0.4 g) was added. The mixture was heated to 50°C for  
five minutes and the solvent removed by evaporation.  
The residue was treated with anhydrous ethyl acetate  
(10 ml) and then with anhydrous dichloromethane and was  
then cooled in ice. 3,3',5,5'-Tetra-iodo-L-thyronine  
25 N-carboxyanhydride (LT<sub>4</sub>-NCA) was collected by  
filtration, washed with anhydrous hexane and dried in  
vacuo without heating. Yield 0.7 g.

Example 20

A solution of 3,3',5,5'-tetraiodo-L-thyronine  
30 N-carboxyanhydride (1.0 g) in anhydrous dioxane (10 ml)  
was stirred rapidly and a 1% solution of sodium

- 25 -

methoxide in methanol (0.05 ml) was added. The mixture was stirred for four days at ambient temperature and then water (10 ml) was added with vigorous stirring. Polymeric 3,3',5,5'-tetraiodo-L- thyronine (poly-LT<sub>4</sub>) was collected by filtration and washed with water. The washed solid was frozen on dry ice and lyophilised under vacuum (less than 0.3 mm Hg) using a shelf temperature of 35°C. Yield 0.21 g.

#### Example 21

The following preparation is suitable for oral administration for the treatment of thyroid hormone deficiency:

	Component	Amount
15	1 Iodothyronine polymer	10 to 300 µg (dose dependent)
	2 Corn Starch	30 mg
	3 Lactose	61 mg
	4 Polyvinylpyrrolidone (PVP)	4 mg
20	5 Talcum	5 mg
	6 Sodium ascorbate	5 mg (antioxidant)

The finely powdered iodothyronine polymer is blended to uniformity with corn starch, lactose, PVP and ascorbate, mixed into an aqueous paste with 1.0 ml H<sub>2</sub>O and freeze dried (30° shelf temperature, 0.03 mm Hg). The resulting powder is mixed uniformly with talcum and pressed into tablets.

Example 22

Hypothyroid male rats which were 2 months old and had an average weight of 95 g were prepared by surgical removal of the thyroid gland. Six to eight weeks were allowed for clearance of endogenous thyroid hormones and rats were bled from the tail vein and serum levels of  $LT_3$  and  $LT_4$  were measured by radioimmunoassay (RIA). Rats having high levels of  $LT_4$  due to inadequate thyroidectomy were removed from the study. A control, untreated group of four thyroidectomized rats was used to test for thyroid remnant regeneration during the course of the experiment. The treatment group of six rats received 10  $\mu$ g/day of the iodothyronine polymer (poly- $T_4/T_3$ ) prepared in Example 9 slurried in corn syrup orally by gavage. After 8 days of treatment, serum levels of  $LT_3$  and  $LT_4$  were again measured by RIA. The results obtained are shown below in which the amounts of  $LT_4$  and  $LT_3$  are shown in nanograms per decilitre of blood and the values quoted are the mean values for the groups of animals.

		<u>Before Treatment</u>		<u>After Treatment</u>	
		$LT_3$ (ng/dl)	$LT_4$ ( $\mu$ g/dl)	$LT_3$ (ng/dl)	$LT_4$ ( $\mu$ g/dl)
25	Control	22	0.80	34	0.88
	Treatment	26	<0.28	284	16.7

Since the establishment of adequate blood levels of thyroid hormones is a necessary and sufficient prerequisite for the euthyroid condition, we concluded

- 27 -

that these data demonstrate that the oral administration of the copolymer of  $LT_4$  and  $LT_3$  can treat thyroid hormone deficiencies resulting from insufficient thyroidal release of thyroid hormones.

5    Example 23

          The sodium salt of 3,3',5,5'-tetraiodo-L-thyronine was suspended in water and dilute (1N) hydrochloric acid was added to give a pH of between 4 and 5. The mixture was shaken for 5 minutes and the solid formed  
10    was collected by filtration, washed with water and dried under vacuo at a temperature in the range 30-50°C.    The solid was 3,3',5,5'-tetraiodo-L-thyronine.

          3,3',5,5'-Tetraiodo-L-thyronine (46.1 g) was  
15    suspended in tetrahydrofuran (400 ml) in a metal foil-wrapped vessel. The mixture was heated at 50-55°C under slightly reduced pressure and solvent (100 ml) was removed by distillation. A further portion of tetrahydrofuran (100 ml) was added and the distillation  
20    repeated to collect a total of 200 ml of solvent. A solution of hexachlorodimethylcarbonate (12 g) in tetrahydrofuran (35 ml) was added at 55°C over a period of twenty minutes and the mixture heated at 55°C for 2 hours. A further portion of hexachlorodimethyl-  
25    carbonate (6 g) in tetrahydrofuran (15 ml) was added over ten minutes and the mixture heated at 55°C for 1.5 hours. The mixture was added to dry hexane (3.8 l) at ambient temperature over 20 minutes. A yellow solid was collected by filtration, washed with hexane and  
30    dried in vacuo at 42°C. Yield 42.7 g.

          A sample (35 g) of the dried product was dissolved in a mixture of ethyl acetate (240 ml) and tetrahydrofuran (90 ml) with warming. The mixture was filtered

- 28 -

through charcoal and the tetrahydrofuran removed by distillation under reduced pressure. Additional ethyl acetate (100 ml) was added and 100 ml of solvent was removed by distillation under reduced pressure. This addition/distillation cycle was repeated twice. The volume was then reduced to 50 ml by evaporation and the solution stored at 2°C for two hours. The resulting solid was collected by filtration, washed with ethyl acetate and dried in vacuo at 40°C. The dried solid was dissolved in a mixture of ethyl acetate (170 ml) and tetrahydrofuran (90 ml). The solvent was removed by evaporation. The residue was dissolved in ethyl acetate (100 ml) and the solvent removed by evaporation. This dissolution/evaporation cycle was repeated and the residue dissolved in ethyl acetate (100 ml). The volume was reduced to 50 ml by evaporation and the solution stored at 2°C for two hours. The resulting solid was collected by filtration, washed with ethyl acetate and dried in vacuo at 40°C to give 3,3',5,5'-tetraiodo-L-thyronine N-carboxyanhydride. Yield 21.2 g.

A solution of 3,3',5,5'-tetraiodo-L-thyronine N-carboxyanhydride (1 g) in dioxane (8 ml) was stirred at 50°C in a metal-foil wrapped vessel. A portion (50 microlitres) of a solution of sodium methoxide (67.2 mg) in methanol (10 ml) was added and the mixture stirred at 50°C for 3.5 hours and then a further portion (50 microlitres) of the above sodium methoxide solution was added and the mixture stirred at 50°C for a total of 23 hours. A solid was collected by filtration, washed with dioxane and dried in vacuo at 50°C to give a dioxane-insoluble fraction of polymeric 3,3',5,5'-tetraiodo-L-thyronine. Yield 0.317 g. The filtrate was slowly added to hexane (25 ml) and the resulting solid collected by filtration, washed with hexane and dried in vacuo at 50°C to give a dioxane-

- 29 -

soluble fraction of polymeric 3,3',5,5'-tetraiodo-L-thyronine. Yield 0.403 g.

Examples 24 to 27

Four solutions of 3,3',5,5'-tetraiodo-L-thyronine N-carboxyanhydride (1 g prepared in Example 23) in dioxane (10 ml) were prepared with warming. Four solutions of sodium methoxide in methanol were made up as follows:-

Ex	Wt of NaOMe (mg)	Volume of methanol (ml)
10		
24	67.2	10
25	134	10
26	335	10
15	27	670

A sample (50 microlitres) of one of these sodium methoxide solutions was added to each of the solutions of 3,3',5,5'-tetraiodo-L-thyronine N-carboxyanhydride and the resulting mixtures were stored at 48-50°C for 23 hours. In Example 24 a further portion (50 microlitres) of the sodium methoxide solution was added after 1 hour. The mixtures were then stored at ambient temperature for 48 hours and the solid which had been formed was collected by filtration and dried at 40°C in vacuo to give dioxane-insoluble fractions of polymeric 3,3',5,5'-tetraiodo-L-thyronine. The yields were Ex 24 0.014 g, Ex 25 0.220 g, Ex 26 0.260 g and Ex 27 0.290 g. The filtrates were added dropwise to hexane (15 ml) over 5 minutes. A precipitate formed which was allowed to settle. The supernatant was removed by decantation and the residue dried in vacuo at 40°C to give dioxane-soluble fractions of polymeric

- 30 -

3,3',5,5'-tetraiodo-L-thyronine as a yellow solid. Yield Ex 24 0.705 g, Ex 25 0.700 g, Ex 26 0.500 g and Ex 27 0.580 g.

#### Example 28

5 A portion (10 microlitres) of a solution of sodium methoxide (134 mg) in methanol (2 ml) was added to a solution of 3,3',5,5'-tetraiodo-L-thyronine N-carboxy-anhydride (1 g prepared in Example 23) in dioxane (10 ml) and the mixture heated at 50°C for 24 hours.  
10 The reaction mixture was centrifuged (3000 rpm for 5 minutes) and the supernatant removed by decantation. The residue was treated with dioxane (8 ml) and the resulting mixture centrifuged (3000 rpm for 5 minutes), and the residue was dried in vacuo at 38°C to give  
15 polymeric 3,3',5,5'-tetraiodo-L-thyronine. Yield 0.485 g.

#### Example 29

A portion (10 microlitres) of a 16% solution of N,N-diisopropylethylamine in dioxane was added to a  
20 solution of 3,3',5,5'-tetraiodo-L-thyronine N-carboxy-anhydride (1 g prepared in Example 23) in dioxane (10 ml) and the mixture heated at 50°C for 96 hours and then centrifuged (3000 rpm for 5 minutes). The supernatant was removed by decantation and the residue  
25 washed with dioxane (4 ml) and dried in vacuo at ambient temperature to give polymeric 3,3',5,5'-tetraiodo-L-thyronine. Yield 0.56 g.

#### Example 30

A portion (5 microlitres) of a 17.2% w/v solution  
30 of sodium methoxide in methanol was added to a solution of 3,3',5,5'-tetraiodo-L-thyronine N-carboxy-anhydride



- 31 -

(1 g prepared in Example 23) in dioxane (10 ml) and the mixture heated at 50°C for 20 hours. The volume of the reaction mixture was reduced to 3 to 4 ml and then the mixture was centrifuged (3000 rpm for 5 minutes) and the supernatant removed by decantation. The residue was washed with dioxane (1 ml) and water (2 x 2 ml) and then slurried with water (10 ml) and then cooled to -78°C and dried in vacuo (0.1 mm Hg) to give polymeric 3,3',5,5'-tetraiodo-L-thyronine. Yield 0.550 g.


#### Example 31

A portion (5 microlitres) of a 17/2% w/v solution of sodium methoxide in methanol was added to a solution of 3,3',5,5'-tetraiodo-L-thyronine N-carboxyanhydride (1 g prepared in Example 23) in tetrahydrofuran (10 ml) and the mixture heated at 48°C for 15 hours. Water was added to precipitate a solid which was separated by decantation. A portion of the wet solid (XX g) was cooled to -78°C and dried in vacuo (0.1 mm Hg) to give polymeric 3,3',5,5'-tetra-iodo-L-thyronine. Yield 0.300 g.

- 32 -

Claims


1. A substantially pure synthetic iodothyronine polymer having a plurality of recurring units, which may be the same or different, of the formula I



- 5 in which A is iodo and B, C and D are independently H or iodo.
2. An iodothyronine polymer as claimed in claim 1 in which polymer substantially all of the recurring units of formula I are in the same stereoisomeric form.
- 10 3. An iodothyronine polymer as claimed in claim 2 in which substantially all of the recurring units of formula I are L-stereoisomers.
4. An iodothyronine polymer as claimed in any one of the preceding claims in which A and C are iodo and B and D are independently H or iodo.
- 15 5. An iodothyronine polymer as claimed in claim 4 in which at least one of B and D is iodo.

6. An iodothyronine polymer as claimed in any one of the preceding claims in which the number of recurring units in the polymer is in the range 5 to 400.
7. An iodothyronine polymer as claimed in any one of  
5 claims 1 to 5 in which the number of recurring units in the polymer is in the range 10 to 400.
8. An iodothyronine polymer as claimed in any one of the preceding claims in which B, C and D are iodo in substantially all of the recurring units.
- 10 9. An iodothyronine polymer as claimed in any one of claims 1 to 7 comprising from 70 to 90% of a recurring unit of formula I in which A, B, C and D are iodo and from 10 to 30% of a recurring unit of formula I in which A and C are iodo, B and D are independently H or  
15 iodo provided that at least one of B or D is H.
10. An iodothyronine polymer as claimed in any one of claims 1 to 7 which has further recurring units of formula II
- $$\text{---} \left[ \text{NH} - \overset{\text{R}}{\underset{|}{\text{CH}}} - \text{CO} \right] \text{---} \quad \text{II}$$
- 20
- in which R is a residue of any of the twenty alpha amino acids commonly found in nature.
11. An iodothyronine polymer as claimed in claim 10 in which the units of formula II are derived from one or  
25 more amino acids selected from the group consisting of lysine, arginine, aspartic acid, glutamic acid, serine or threonine.
12. An N-carboxyanhydride compound of formula III

- 34 -




in which A is iodo and B, C and D are independently H or iodo provided that at least one of B, C or D is H.

13. An L-stereoisomer of a N-carboxyanhydride compound of formula III in which A is iodo and B, C and D are independently H or iodo.

14. An N-carboxyanhydride compound as claimed in claim 13 in which A and C are iodo and at least one of B and D is iodo.

15. A pharmaceutical composition suitable for treating thyroid hormone deficiencies comprising a pharmaceutically acceptable diluent or carrier and a pharmacologically active ingredient consisting of a pharmaceutically effective amount of a substantially pure synthetic iodothyronine polymer having a plurality of recurring units, which may be the same or different, of the formula I

- 35 -



in which A and C are iodo and B and D are independently H or iodo, in which polymer substantially all of the recurring units of formula I are L-stereoisomers.

16. A pharmaceutical composition as claimed in claim 5 15 in which at least one of B or D is iodo.


17. A pharmaceutical composition as claimed in claim 15 or 16 in which the number of recurring units of formula I in the iodothyronine polymer is in the range 5 to 400.

10 18. A pharmaceutical composition as claimed in any one of the preceding claims in which B, C and D are iodo in substantially all of the recurring units.

15 19. A pharmaceutical composition as claimed in any one of claims 15 to 17 in which the iodothyronine polymer comprises from 70 to 90% of a recurring unit of formula I in which A, B, C and D are iodo and from 10 to 30% of a recurring unit of formula I in which A and C are iodo, B and D are independently H or iodo provided that at least one of B or D is H.

- 36 -

20. A pharmaceutical composition as claimed in any one of claims 15 to 18 in which the iodothyronine polymer has further recurring units of formula II



5 in which R is a residue of any one of the twenty amino acids commonly found in nature.

21. A pharmaceutical composition as claimed in claim 20 in which the further recurring units of formula II are derived from one or more amino acids selected from the group consisting of lysine, arginine, aspartic  
10 acid, glutamic acid, serine or threonine.

22. A pharmaceutical composition as claimed in claim 15 in which the pharmacologically active ingredient comprises a mixture of pharmaceutically effective amounts of a first iodothyronine polymer in which B, C and D are iodo and a second iodothyronine polymer in which B and C are iodo and D is H.


23. A pharmaceutical composition as claimed in claim 22 in which the molar ratio of the amount of the first iodothyronine polymer to the amount of the second  
20 iodothyronine polymer lies in the range 7:3 to 9:1.

24. A pharmaceutical composition as claimed in claim 22 which also comprises a pharmaceutically effective amount of a further iodothyronine polymer in which B is H and C and D are iodo.

- 37 -

25. A pharmaceutical composition as claimed in claim 24 in which the ratio of the amount of the first iodothyronine polymer to the amount of the second iodothyronine polymer to the amount of the further  
5 iodothyronine polymer is apporoximately 80:15:5.

26. A method of treating thyroid hormone deficiencies in a patient in need of such treatment which consists essentially of administering to a patient in need thereof a pharmaceutically effective amount of a  
10 synthetic iodothyronine polymer having a plurality of recurring units, which may be the same or different, of the formula I



in which A and C are iodo and B and D are independently H or iodo, in which polymer substantially all of the  
15 recurring units of formula I are L-stereoisomers.

27. A method of treating thyroid hormone deficiencies as claimed in claim 26 in which at least one of B or D is iodo.

28. A method of treating thyroid hormone deficiencies  
20 as claimed in claim 26 or 27 in which the number of


- 38 -

recurring units of formula I in the iodothyronine polymer is in the range 5 to 400.

29. A method of treating thyroid hormone deficiencies as claimed in any one of claims 26 to 28 in which B, C and D are iodo in substantially all of the recurring units.

30. A method of treating thyroid hormone deficiencies as claimed in any one of claims 26 to 28 in which the iodothyronine polymer comprises from 70 to 90% of a recurring unit of formula I in which A, B, C and D are iodo and from 10 to 30% of a recurring unit of formula I in which A and C are I, B and D are independently H or iodo provided that at least one of B or D is H.

31. A method of treating thyroid hormone deficiencies as claimed in any one of claims 26 to 30 in which the iodothyronine polymer has further recurring units of formula II




in which R is a residue of any one of the twenty amino acids commonly found in nature.

32. A method of treating thyroid hormone deficiencies as claimed in claim 31 in which the further recurring units of formula II are derived from one or more amino acids selected from the group of lysine, arginine, aspartic acid, glutamic acid, serine or threonine.

33. The use of a synthetic iodothyronine polymer having a plurality of recurring units, which may be the same or different, of the formula I




- 39 -



in which A and C are iodo and B and D are independently H or iodo in which polymer substantially all of the recurring units of formula I are L-stereoisomers in the treatment of thyroid hormone deficiencies.

- 5 34. The use of a synthetic iodothyronine polymer having a plurality of recurring units, which may be the same or different, of the formula I




in which A and C are iodo and B and D are independently H or iodo in which polymer substantially all of the recurring units of formula I are L-stereoisomers in the

10

- 40 -

manufacture of a medicament for the treatment of thyroid hormone deficiencies.

35. A process for preparing an iodothyronine polymer having recurring units of formula I in which A is iodo and B, C and D are independently H or iodo comprising the steps of polymerising one or more N-carboxyanhydrides of formula III



- in which A is iodo and B, C and D are independently H or iodo to produce the polymer and then recovering said polymer in substantially pure form.

36. A process as claimed in claim 35 in which the polymerisation step comprises reacting one or more N-carboxyanhydrides in an anhydrous solvent using a base as initiator.


37. A process as claimed in claim 35 for preparing an iodothyronine polymer comprising from 70 to 90% of a recurring unit of formula I in which A, B, C and D are

- 41 -

iodo and from 10 to 30% of a recurring unit of formula I in which A, B and C are iodo and D is H, said process comprising the steps of polymerising a mixture of 70 to 90% of an N-carboxyanhydride of formula III in which A, B, C and D are iodo and 30 to 10% of an N-carboxyanhydride of formula III in which A, B and C is iodo and D is H and then recovering said polymer in substantially pure form.

38. A process as claimed in claim 37 in which the polymerisation step comprises reacting one or more N-carboxyanhydrides in an anhydrous solvent using a base as initiator.

39. A process for preparing an iodothyronine polymer which contains recurring units of formula I and further recurring units of formula II which comprises the step of copolymerising an N-carboxyanhydride of formula III and an N-carboxyanhydride of formula IV




in which R is a residue of any of the twenty alpha amino acids commonly found in nature, then recovering the iodothyronine polymer in substantially pure form.

40. A process as claimed in claim 39 in which the copolymerisation step comprises reacting the N-carboxyanhydrides in an anhydrous solvent using a base as initiator.

- 42 -

41. A process for preparing an iodothyronine polymer having recurring units of formula I in which A is iodo, B is H or iodo and C and D are iodo comprising the steps of diiodinating an iodothyronine polymer having recurring units of formula I in which A is iodo, B is H or iodo and C and D are H and then recovering the required iodothyronine polymer in substantially pure form.
42. A process as claimed in claim 41 in which the diiodination step comprises the use of iodine monochloride as the diiodinating reagent.
43. A process for preparing an iodothyronine polymer having recurring units of formula I in which A is iodo, B is H or iodo, C is iodo and D is H comprising the steps of monoiodinating an iodothyronine polymer having recurring units of formula I in which A is iodo, B is H or iodo and C and D are H and then recovering the desired iodothyronine polymer in substantially pure form.
44. A process as claimed in claim 43 in which the iodination step comprises the use of potassium triiodide as the monoiodinating agent.
45. A process for preparing an N-carboxyanhydride of formula III comprises the step of reacting an iodothyronine of formula V

- 43 -




V

or a salt thereof with a carbonylating agent.

46. A process as claimed in claim 45 in which the carbonylating agent is hexachlorodimethylcarbonate or phosgene.

# INTERNATIONAL SEARCH REPORT

International Application No PCT/US 90/06213

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC5: C 07 K 15/00, C 08 G 69/10, C 07 K 13/00, C 07 K 7/00 A 61 K 37/02, A 61 K 37/24, C 07 D 263/44		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
IPC5	C 07 K; C 08 G, A 61 K; C 07 D	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category *	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
A	J. Am. Chem. Soc., Vol. 54, September 1932, Charles S. Myers: "Some derivatives of diiodotyrosine and thyroxine. The action of acetic anhydride on diiodotyrosine ", see page 3718 - page 3725  --	12-14
A	The Peptides, Vol. 1, 1965, Eberhard Schröder et al.: "Methods of Peptide Synthesis ", see page 122 - page 124  --	12-14, 35-38
A	DE, A1, 2416941 (HELMUT PRATZEL) 9 October 1975, see pages 1-5  --	12-14, 35-38
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents: <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p> </div> </div>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
14th February 1991	05.03.91	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	 p.p. MISS D. S. KOWALCZYK	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
A	Journal of Polymer Science: Polymer Chemistry Edition, Vol. 21, 1983, Brent A. Burdick: "Preparation and Polymerization of Thyroxine Methacrylate Monomers ", see page 1997 - page 2001  --	1
A	EP, A1, 0327411 (DELALANDE S.A.) 9 August 1989, see the whole document  --  -----	45,46

## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE <sup>1</sup>

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers 26-33 because they relate to subject matter not required to be searched by this Authority, namely:

See PCT Rule 39.1(iv)

Method for treatment of the human or animal body by means of surgery or therapy, as well as diagnostic methods.

2. ☐ Claim numbers ..... because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers ..... because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING <sup>2</sup>

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

## Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO. PCT/US 90/06213**

SA 41942

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.  
The members are as contained in the European Patent Office EDP file on 28/12/90  
The European Patent office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE-A1- 2416941	09/10/75	FR-A- 2243183 US-A- 3951741	04/04/75 20/04/76
EP-A1- 0327411	09/08/89	FR-A-B- 2625507 JP-A- 1210425	07/07/89 24/08/89

For more details about this annex : see Official Journal of the European patent Office, No. 12/82